

MULTIPLE AUTOIMMUNE REACTIVITY SCREEN™



2 mL serum

- ▶ Parietal Cell + ATPase IgG + IgA Combined
- ▶ Intrinsic Factor IgG + IgA Combined
- ▶ ASCA + ANCA IgG + IgA Combined
- ▶ Tropomyosin IgG + IgA Combined
- ▶ Thyroglobulin IgG + IgA Combined
- ▶ Thyroid Peroxidase (TPO) IgG + IgA Combined
- ▶ 21 Hydroxylase (Adrenal Cortex) IgG + IgA Combined
- ▶ Myocardial Peptide IgG + IgA Combined
- ▶ Alpha-Myosin IgG + IgA Combined
- ▶ Phospholipid IgG + IgA Combined
- ▶ Platelet Glycoprotein IgG + IgA Combined
- ▶ Ovary/Testis* IgG + IgA Combined
- ▶ Fibulin IgG + IgA Combined
- ▶ Collagen Complex IgG + IgA Combined
- ▶ Arthritic Peptide IgG + IgA Combined
- ▶ Osteocyte IgG + IgA Combined
- ▶ Cytochrome P450 (Hepatocyte) IgG + IgA Combined
- ▶ Insulin + Islet Cell Antigen IgG + IgA Combined
- ▶ Glutamic Acid Decarboxylase 65 (GAD 65) IgG + IgA Combined
- ▶ Myelin Basic Protein IgG + IgA Combined
- ▶ Asialoganglioside IgG + IgA Combined
- ▶ Alpha + Beta Tubulin IgG + IgA Combined
- ▶ Cerebellar IgG + IgA Combined
- ▶ Synapsin IgG + IgA Combined

* Ovary and Testis are tested together to avoid any confusion arising out of potential cross-reactivity.

CLINICAL USE:

- Measure Predictive Antibodies, some of which can appear up to ten years before the clinical onset of disease
- Economically and efficiently assess possible tissue damage to multiple organs of the body

RECOMMENDED FOR PATIENTS WITH:

- A diagnosed autoimmune disorder, as they commonly develop additional autoimmune conditions
- Chronic increased intestinal permeability, which is the gateway for environmentally-induced autoimmune disorders



www.CyrexLabs.com

MULTIPLE AUTOIMMUNE REACTIVITY SCREEN™

